MycoKeys

MycoKeys 100: 153–170 (2023)
DOI: 10.3897/mycokeys.100.112458

Research Article

Exploring diversity within the genus *Tulostoma* (Basidiomycota, Agaricales) in the Pannonian sandy steppe: four fascinating novel species from Hungary

Péter Finy[1,2], Mikael Jeppson[3], Dániel G. Knapp[1,4], Viktor Papp[5], László Albert[2], István Ölvedi[2], Károly Bóka[1], Dóra Varga[1], Gábor M. Kovács[1], Bálint Dima[1]

1 *Department of Plant Anatomy, Institute of Biology, Eötvös Loránd University, Pázmány Péter sétány 1/C, Budapest 1117, Hungary*
2 *Hungarian Mycological Society, Könyves Kálmán krt. 40. Budapest 1087, Hungary*
3 *University of Gothenburg, Biological and Environmental Sciences, P.O. Box 461, SE-40530 Göteborg, Sweden*
4 *Department of Forestry and Wood Technology, Linnaeus University, Växjö, Sweden*
5 *Department of Botany, Hungarian University of Agriculture and Life Sciences, Villányi út 29–43, H-1118 Budapest, Hungary*
Corresponding author: Bálint Dima (cortinarius1@gmail.com)

Abstract

Steppe vegetation on sandy soil in Hungary has recently been revealed as one of the hot spots in Europe for the stalked puffballs (genus *Tulostoma*). In the framework of the taxonomic revision of gasteroid fungi in Hungary, four *Tulostoma* species are described here as new to science: *T. dunense*, *T. hungaricum*, *T. sacchariolens* and *T. shaihuludii*. The study is based on detailed macro- and micromorphological investigations (including light and scanning electron microscopy), as well as a three-locus phylogeny of nrDNA ITS, nrDNA LSU and *tef1-α* sequences. The ITS and LSU sequences generated from the type specimen of *T. cretaceum* are provided and this resolved partly the taxonomy of the difficult species complex of *T.* aff. *cretaceum*.

Key words: Gasteroid, hot spot, molecular systematics, Pannonian inland sand dune thicket, phylogeny, taxonomy, Tulostomataceae

OPEN ACCESS

Academic editor: María P. Martín
Received: 9 September 2023
Accepted: 27 October 2023
Published: 29 November 2023

Citation: Finy P, Jeppson M, Knapp DG, Papp V, Albert L, Ölvedi I, Bóka K, Varga D, Kovács GM, Dima B (2023) Exploring diversity within the genus *Tulostoma* (Basidiomycota, Agaricales) in the Pannonian sandy steppe: four fascinating novel species from Hungary. MycoKeys 100: 153–170. https://doi.org/10.3897/mycokeys.100.112458

Copyright: © Péter Finy et al.
This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

Introduction

The genus *Tulostoma* was erected by Persoon (1794, 1801) encompassing two species, *T. brumale* and *T. squamosum*. Several new species have since been added from all continents, except Antarctica. In a monograph of the genus based on type studies, Wright (1987) accepted 139 species worldwide. Later studies generally confirmed those species concepts, nevertheless reduced some of the species to synonymy (e.g. Moreno et al. (1992, 1997); Altés et al. (1999); Jeppson et al. (2017)). With the introduction of molecular methods in taxonomy, unexpected species diversity has been detected and new, formerly unknown species have been described. In Europe, Jeppson et al. (2017) suggested Mediterranean grassland regions of the Iberian Peninsula, as well as steppe habitats in East Central Europe, as hot-spot areas for species diversity in *Tulostoma*. Jeppson et al. (2017)

described two novel species with type localities in Central Hungary (*T. grandisporum*, *T. pannonicum*), but their phylogenetic and morphological results indicated the presence of at least 19 previously-undescribed European species of *Tulostoma*, nine of which had been collected in Hungary. The species diversity in Eastern Europe was further emphasised by Rusevska et al. (2019) who reported four species from North Macedonia distinct from all known and described species.

In Europe, Hungary has an exceptionally large diversity of gasteroid taxa mainly due to the suitable habitats of the Pannonian sandy steppe areas of the country (Fig. 1). The *Festucetum vaginatae* plant communities are characteristic on open, continental sandy soils, dominated by the grass species *Festuca vaginata* which also occur on open steppe mosaics between the poplar–juniper sand dune thickets (Bölöni et al. 2011; Rimóczi et al. 2011). Gasteroid fungi occur especially in those areas where *Stipa borysthenica*, *Fumana procumbens* or *Juniperus communis* are present (Fig. 1).

In this paper, we propose four species of *Tulostoma* new to science, two of which were retrieved previously by Jeppson et al. (2017) as *Tulostoma* sp. 1 and *T.* aff. *cretaceum*. Additionally, we present two further species identified through subsequent collections and field investigations. One of the new species was previously reported from Hungary under the name *T. volvulatum* (Hollós 1904; Siller and Vasas 1995; Siller et al. 2005) and *T. obesum* (Siller et al. 2006; Rimóczi et al. 2011) which is listed as a protected species by Hungarian law.

Figure 1. Habitats of *Tulostoma* species in Hungary: **a** *T. hungaricum* in Orgovány **b** *T. sacchariolens* in Orgovány **c** *T. dunense* in Izsák (Soltszentimre) **d** *T. shaihuludii* in Izsák (Soltszentimre). Photos: P. Finy.

Materials and methods

Samples of *Tulostoma* were collected in Hungary over a period of more than 25 years. Collecting has mostly been performed in the sandy habitats of the Great Hungarian Plain on both sides of the Danube (Kiskunság, Mezőföld). Studied collections were deposited in the herbaria BP (only holotypes), GB and in the Department of Plant Anatomy, Eötvös Loránd University (abbreviated further as ELTE).

Macromorphological study

Mature basidiomata of *Tulostoma* were collected and studied under a stereomicroscope, regarding their macromorphological characteristics (size, colour, shape of the spore-sac (capitulum), type of mouth (ostiole), type of exoperidium as well as features of the stem), in accordance with Wright (1987). In situ or ex situ photo-documentation of each sample was carried out.

Microscopy

Microscopic features were studied under an Olympus BH-2 light microscope. Samples were mounted in lactophenol-cotton blue and heated to boiling temperature. Measurements were performed under 1000× magnification and calculated digitally using Piximètre software (www.piximetre.fr). Spore dimensions are given without the ornamentation of the spore walls. Small pieces of peridium and gleba from dried basidiomata were prepared, fixed to stubs, coated with gold and examined under a Hitachi S2460N (Hitachi Ltd., Tokyo, Japan) scanning electron microscope (SEM) at 22 kV accelerating voltage.

Molecular study

Total DNA extraction was carried out with the E.Z.N.A. SP Fungal DNA Mini Kit (Omega Bio-Tek, Norcross, GA, USA) and NucleoSpin Plant II Mini Kit (Macherey-Nagel, Düren, Germany) following the instructions of the manufacturers. The ITS (internal transcribed spacer) region of the nrDNA which is the universal fungal DNA barcode region (Schoch et al. 2012) was amplified using the primer pairs ITS1F/ITS4 (White et al. 1990; Gardes and Bruns 1993) as described in Papp and Dima (2018). The primers LR0R (Rehner and Samuels 1994) and LR5 (Vilgalys and Hester 1990) were used to amplify the partial 28S nrRNA gene (LSU) of the nrDNA operon region. The primers EF1-983F and EF1-2218R (Rehner and Buckley 2005) were used to amplify part of the translation elongation factor 1α (*tef1-α*). Sequencing of the amplicons with the primers used for amplification was carried out by LGC Genomics (Berlin, Germany). The sequences were compiled from electropherograms using the Staden software package (Staden et al. 2000) and CodonCode Aligner package (CodonCode Corp., Centerville, Massachusetts, USA). Sequences of each locus (ITS, LSU and *tef1-α*), together with sequences of respective species downloaded from GenBank mainly based on Jeppson et al. (2017), were aligned separately with the online MAFFT version 7 using the E-INS-i strategy (Katoh and Standley 2013) (Table 1). The alignments were checked and edited in MEGA7 (Kumar et al. 2016) and concatenated to one dataset in SeaView 5 (Gouy et al. 2021). Bayesian Inference (BI) analyses were performed with MrBayes 3.1.2 (Ron-

quist and Huelsenbeck 2003) using a GTR + G substitution model. Four Markov chains were run for 10,000,000 generations, sampling every 1,000 generations with a burn-in value set at 4,000 sampled trees. Maximum Likelihood (ML) phylogenetic analysis was carried out with the raxmlGUI 1.3 implementation (Silvestro and Michalak 2012; Stamatakis 2014). The GTR + G nucleotide substitution model and ML estimation of base frequencies were applied for the partitions. ML bootstrap (BS) analysis with 1,000 replicates was used to test the support of the branches. *Tulostoma pulchellum* (MJ7833) and *T. striatum* (Fritz 2010-2) served as outgroups. Intra- and interspecific genetic differences were calculated by dividing the number of differences (substitutions and/or indels) found in the whole ITS region by the length of the region. Phylogenetic trees were visualised and edited in MEGA 7 (Kumar et al. 2016) and deposited together with the alignments at Figshare repository (10.6084/m9.figshare.24112749). Newly-generated sequences were submitted to GenBank. Studied voucher collections are presented in Table 1.

Table 1. Sequences used in this study. Newly-generated sequences are marked in boldface.

Name	Strain/Voucher	Country	ITS	LSU	TEF	References
Tulostoma ahmadii	HUP SH-33b, holotype	Pakistan	KP738712	–	–	Hussain et al. (2016)
Tulostoma albicans	B2092, P.S. Catcheside 1266	Australia	–	MK278628	–	Varga et al. (2019)
Tulostoma albicans	Cope, NY, Holotype	United States	KX576548	–	–	Jeppson et al. (2017)
Tulostoma beccarianum	Finy2	Hungary	KU519076	KU519076	KU843959	Jeppson et al. (2017)
Tulostoma beccarianum	Herb. Bresadola (S), holotype	Italy	KX640979	–	–	Jeppson et al. (2017)
Tulostoma berkeleyi	JLH MyCoPortal 6604754	United States	MK578704	MK578704	–	Unpublished
Tulostoma brumale	Finy9	Hungary	KU519059	KU519059	KU843944	Jeppson et al. (2017)
Tulostoma calcareum	Finy4	Hungary	KU519088	KU519088	KU843895	Jeppson et al. (2017)
Tulostoma calcareum	MJ6965, holotype	Sweden	KU519086	KU519086	KU843881	Jeppson et al. (2017)
Tulostoma calongei	MJ8773, holotype	Spain	KU518973	KU518973	KU844000	Jeppson et al. (2017)
Tulostoma caespitosum cf.	SNMH9	Slovakia	MK907419	MK907419	–	Unpublished
Tulostoma caespitosum cf.	MJ881114	Spain	KU519031	KU519031	KU843978	Jeppson et al. (2017)
Tulostoma caespitosum cf.	AH15040	Spain	KU519032	KU519032	KU843979	Jeppson et al. (2017)
Tulostoma cretaceum	**NY737977, holotype**	**United States**	**OR722641**	**OR722660**	–	**This study**
Tulostoma cretaceum cf. 1	Knudsen0107	Russia	KU518993	KU518993	KU843988	Jeppson et al. (2017)
Tulostoma cretaceum cf. 2	AH13672	Spain	KU518998	KU518998	KU843991	Jeppson et al. (2017)
Tulostoma cretaceum cf. 2	AH3995	Spain	KU518999	KU518999	KU843992	Jeppson et al. (2017)
Tulostoma cretaceum cf. 2	MJ6194	Spain	KU518997	KU518997	KU843989	Jeppson et al. (2017)
Tulostoma cretaceum cf. 2	MJ9304	Spain	KU519000	KU519000	KU843990	Jeppson et al. (2017)
***Tulostoma cretaceum* cf. 3**	**FP-2023-05-11-1**	**Kazakhstan**	**OR722639**	**OR722658**	**–**	**This study**
***Tulostoma cretaceum* cf. 3**	**FP-2023-05-11-4**	**Kazakhstan**	**OR722640**	**OR722659**	**–**	**This study**
Tulostoma cretaceum cf. 3	SNMH10	Kazakhstan	MK907420	MK907420	–	Unpublished
Tulostoma cretaceum cf.	MJ3821	Hungary	KU518994	KU518994	KU843993	Jeppson et al. (2017)
Tulostoma cyclophorum	MJ8862	Hungary	KU518985	KU518985	KU843963	Jeppson et al. (2017)
Tulostoma domingueziae	MLHC24 (CORD), holotype	Argentina	HQ667594	HQ667597	–	Caffot et al. (2011)
Tulostoma dunense	**BP112640, holotype**	**Hungary**	**OR722622**	**OR722648**	**OR707014**	**This study**
Tulostoma dunense	**DB-2021-11-21-2**	**Hungary**	**OR722626**	**–**	**–**	**This study**
Tulostoma dunense	**FP-2019-12-07**	**Hungary**	**OR722617**	**OR722643**	**OR707009**	**This study**
Tulostoma dunense	**FP-2020-12-06**	**Hungary**	**OR722618**	**OR722644**	**OR707010**	**This study**
Tulostoma dunense	**FP-2022-01-02-1**	**Hungary**	**OR722619**	**OR722645**	**OR707011**	**This study**
Tulostoma dunense	**FP-2021-01-02**	**Hungary**	**OR722620**	**OR722646**	**OR707012**	**This study**
Tulostoma dunense	**FP-2016-06-05**	**Hungary**	**OR722621**	**OR722647**	**OR707013**	**This study**
Tulostoma dunense	**FP-2021-02-18**	**Hungary**	**OR722623**	**OR722649**	**OR707015**	**This study**
Tulostoma dunense	**FP-2015-12-06**	**Hungary**	**OR722624**	**OR722650**	**OR707016**	**This study**
Tulostoma dunense	**FP-2016-12-11**	**Hungary**	**OR722625**	**OR722651**	**OR707017**	**This study**
Tulostoma dunense	MJ6103 (as cf. *cretaceum*)	Hungary	KU518995	KU518995	KU843994	Jeppson et al. (2017)
Tulostoma dunense	MJ7759 (as cf. *cretaceum*)	Hungary	KU518996	KU518996	KU843995	Jeppson et al. (2017)

Name	Strain/Voucher	Country	ITS	LSU	TEF	References
Tulostoma eckbladii	Sivertsen930717, TRH9565, holotype	Norway	KU519069	KU519069	KU843952	Jeppson et al. (2017)
Tulostoma excentricum	BPI 729284, holotype	United States	KU519055	KU519055	–	Jeppson et al. (2017)
Tulostoma fimbriatum	Finy8	Hungary	KU518968	KU518968	KU843912	Jeppson et al. (2017)
Tulostoma fimbriatum	Månsson 991010, epitype	Sweden	KU518963	KU518963	KU843904	Jeppson et al. (2017)
Tulostoma fulvellum	Kabát 970428	Slovakia	KU518991	KU518991	KU844001	Jeppson et al. (2017)
Tulostoma giovanellae	MJ8706	Spain	KU519071	KU519071	KU843954	Jeppson et al. (2017)
Tulostoma grandisporum	Finy10	Hungary	KU519005	KU519005	KU843922	Jeppson et al. (2017)
Tulostoma grandisporum	MJ8907, holotype	Hungary	KU519003	KU519003	KU843924	Jeppson et al. (2017)
Tulostoma hungaricum	**BP112641, holotype**	**Hungary**	**OR722630**	**OR722653**	**–**	**This study**
Tulostoma hungaricum	**FP-2019-01-23**	**Hungary**	**OR722627**	**–**	**–**	**This study**
Tulostoma hungaricum	**FP-2021-02-19**	**Hungary**	**OR722628**	**–**	**–**	**This study**
Tulostoma hungaricum	**FP-2022-01-02-2**	**Hungary**	**OR722629**	**OR722652**	**OR707021**	**This study**
Tulostoma kotlabae	Brůžek 140918	Czech Republic	KU519028	KU519028	KU843977	Jeppson et al. (2017)
Tulostoma kotlabae	Kotlaba (PRM 704203), holotype	Slovakia	KX576544	KX576544	–	Jeppson et al. (2017)
Tulostoma cf. *kotlabae*	MJ5996	Hungary	KU519016	KU519016	KU843966	Jeppson et al. (2017)
Tulostoma cf. *kotlabae*	Finy1	Hungary	KU519017	KU519017	KU843967	Jeppson et al. (2017)
Tulostoma cf. *kotlabae*	MJ7795	Hungary	KU519020	KU519020	KU843970	Jeppson et al. (2017)
Tulostoma laceratum	**NY834492**	**United States**	**OR722642**	**OR722661**	–	**This study**
Tulostoma lloydii	Lahti 201210	Italy	KU518990	KU518990	KU843965	Jeppson et al. (2017)
Tulostoma lusitanicum	LISU-MGA-8	Portugal	KX576542	KX576542	–	Jeppson et al. (2017)
Tulostoma lysocephalum	Long 9639, holotype	United States	KU519034	KU519034	–	Jeppson et al. (2017)
Tulostoma melanocyclum	MJ090418	Hungary	KU519106	KU519106	KU843890	Jeppson et al. (2017)
Tulostoma cf. *nanum*	MJ4976	Hungary	KU519036	KU519036	KU843968	Jeppson et al. (2017)
Tulostoma niveum	MJ7692	Sweden	KU519078	KU519078	KU843932	Jeppson et al. (2017)
Tulostoma obesum	Cooke 2715, isotype	United States	KX576541	KX576541	–	Jeppson et al. (2017)
Tulostoma obesum	MJ8695	Spain	KU518986	KU518986	KU843985	Jeppson et al. (2017)
Tulostoma pannonicum	MJ7764, holotype	Hungary	KU519010	KU519010	–	Jeppson et al. (2017)
Tulostoma pannonicum	MJ7803	Hungary	KU519011	KU519011	KU843996	Jeppson et al. (2017)
Tulostoma pseudopulchellum	AH 11603, paratype	Spain	KU519012	KU519012	KU843997	Jeppson et al. (2017)
Tulostoma pseudopulchellum	AH 11605, holotype	Spain	KX513827	KX513827	–	Jeppson et al. (2017)
Tulostoma pulchellum	MJ7833	Hungary	KU518957	KU518957	KU843928	Jeppson et al. (2017)
Tulostoma punctatum	BPI 729033, lectotype	United States	KC333071	KC333071	–	Jeppson et al. (2017)
Tulostoma punctatum	MJ7472	Slovakia	KU518952	KU518952	KU843875	Jeppson et al. (2017)
Tulostoma pygmaeum cf.	Brůžek 131207	Slovakia	KU519041	KU519041	KU843931	Jeppson et al. (2017)
Tulostoma rufum	BPI 704578, holotype	United States	KU519107	KU519107	–	Jeppson et al. (2017)
Tulostoma sacchariolens	**BP112642, holotype**	**Hungary**	**OR722632**	**OR722654**	**OR707020**	**This study**
Tulostoma sacchariolens	**FP-2019-12-06**	**Hungary**	**OR722631**	**–**	**–**	**This study**
Tulostoma sacchariolens	**FP-2021-01-24b**	**Hungary**	**OR722633**	**–**	**–**	**This study**
Tulostoma sacchariolens	**FP-2021-02-18**	**Hungary**	**OR722634**	**OR722655**	**–**	**This study**
Tulostoma shaihuludii	**BP112643, holotype**	**Hungary**	**OR722637**	**OR722657**	**OR707019**	**This study**
Tulostoma shaihuludii	**FP-2020-12-01**	**Hungary**	**OR722635**	**–**	**–**	**This study**
Tulostoma shaihuludii	**FP-2020-12-27**	**Hungary**	**OR722636**	**OR722656**	**OR707018**	**This study**
Tulostoma shaihuludii	**FP-2017-12-09**	**Hungary**	**OR722638**	**–**	**–**	**This study**
Tulostoma shaihuludii	MJ7762	Hungary	KU518979	KU518979	KU843981	Jeppson et al. (2017)
Tulostoma simulans	MJ3844	Hungary	KU519052	KU519052	KU843941	Jeppson et al. (2017)
Tulostoma sp. 10	MJ3813	Hungary	KU519029	KU519029	–	Jeppson et al. (2017)
Tulostoma sp. 14	MJ5004	Spain	KU519039	KU519039	KU843999	Jeppson et al. (2017)
Tulostoma sp. 20	MJ5015	Spain	KU519067	KU519067	KU843950	Jeppson et al. (2017)
Tulostoma sp. 21	AH11698	Spain	KX640986	KX640986	–	Jeppson et al. (2017)
Tulostoma squamosum	Larsson 260-06	France	KU519097	KU519097	KU843892	Jeppson et al. (2017)
Tulostoma striatum	Fritz 2010-2	Mongolia	KU518958	KU518958	KU843929	Jeppson et al. (2017)
Tulostoma submembranaceum	AH15132, holotype	Mexico	KX513826	KX513826	–	Jeppson et al. (2017)
Tulostoma submembranaceum cf.	MJ9296	Spain	KU519014	KU519014	KU843984	Jeppson et al. (2017)
Tulostoma subsquamosum	MJ4945	Hungary	KU519091	KU519091	KU843899	Jeppson et al. (2017)
Tulostoma verrucosum	CCB142	United States	MG663293	MG663293	–	Unpublished
Tulostoma winterhoffii	MJ7761	Hungary	KU518976	KU518976	KU843916	Jeppson et al. (2017)
Tulostoma xerophilum	Long 9688, BPI 802484, holotype	United States	KX576549	–	–	Jeppson et al. (2017)

Results

Phylogenetic analysis

The three-locus molecular phylogenetic analyses of the newly-generated and representative *Tulostoma* sequences were based on 94 ITS, 76 LSU and 60 *tef1-α* (Table 1) and 3321 characters. In this study, 26 ITS, 26 LSU and 13 *tef1-α* sequences were newly gained, including the ITS and LSU sequences of the holotype of *Tulostoma cretaceum* (Table 1). Phylogenetic trees from ML and BI analyses showed congruent topologies and the sequences representing the four new species proposed here formed strongly-supported clades (MLBS/BIPP = 100%/1.00). The best scoring ML tree is shown in Fig. 2.

Taxonomy

Tulostoma dunense **Finy, Jeppson, L. Albert, Ölvedi, Dima & V. Papp, sp. nov.**
MycoBank No: MB 849931
Fig. 3

Holotype. HUNGARY, Tolna, Németkér, open sandy grassland, 18 Oct 2020, P. Finy, I. Ölvedi, FP-2020-10-18 (BP112640, isotype GB). GenBank: ITS OR722622, LSU OR722648, tef1 OR707014.

Etymology. The epithet refers to the continental, open, bare sandy habitat of this species, similar to coastal dunes.

Description. Spore-sac subglobose, depressed-globose, 10–20 mm. Exoperidium hyphal, encrusting sand only at the base of the spore-sac. Endoperidium tough, chalky white or dirty-dingy white, with age becoming greyish, young basidiomata with velvety surface. Mouth prominent, fibrillose-lacerate, irregular sometimes remains unopened for a long time and splits later due to mechanical pressure (wind or trampling). Socket distantly separated from the stem. Stem 35–80 × 1.5–5 mm, initially white, then ochraceous, with age greyish–blackish, longitudinally furrowed, at the base with a volva and a prominent, easily broken pseudorhiza. Gleba ferruginous to brick-red brown, usually scattered on the surface of the spore-sac. Capillitium brown, 2.5–10 µm in diameter with walls 0.7–2.5 µm in diameter, fragile, breaking up at septal levels in 40–350 µm long segments with rounded, not widened ends, rarely branching. Spores subglobose to oval, 4.6–5.2 × 4.0–4.8 µm (av. 4.4 × 4.9 µm), smooth under LM and SEM.

Habitat and distribution. The psammophilous species *Tulostoma dunense* known so far only from sandy areas of the Great Hungarian Plain of Hungary. It occurs on both sides of the Danube (Kiskunság, Dél-Mezőföld), where open dunes appear. It mainly grows solitary, deep in the sand in large, open sandy areas to bare spots.

Notes. *Tulostoma dunense* was previously recorded in Hungary by Hollós (1904), Siller and Vasas (1995), Babos (1999), Siller et al. (2005), Siller et al. (2006) and Rimóczi et al. (2011) under the names of *T. volvulatum*, *T. obesum* and *T.* aff. *cretaceum*. Hollós (1904) included both *T. giovanellae* and *T. dunense* under the name *T. volvulatum* (nom. rej., Altés et al. (1999)) and recorded it in urban places in the City of Kecskemét (now *T. giovanellae*) as well as in sand dunes (now the new species, *T. dunense*). The brownish colour of the capillitium characteristic of specimens from sand dunes and largely absent in those from


FP-2022-01-02-1 I Tulostoma dunense I Hungary
FP-2021-02-18 I Tulostoma dunense I Hungary
FP-2020-12-06 I Tulostoma dunense I Hungary
FP-2021-01-02 I Tulostoma dunense I Hungary
FP-2019-12-07 I Tulostoma dunense I Hungary
FP-2015-12-06 I Tulostoma dunense I Hungary
BP112640 I Tulostoma dunense I Hungary I holotype
MJ6103 I Tulostoma dunense (as cf. cretaceum) I Hungary
MJ7759 I Tulostoma dunense (as cf. cretaceum) I Hungary
FP-2016-06-05 I Tulostoma dunense I Hungary
DB-2021-11-21-2 I Tulostoma dunense I Hungary
FP-2016-12-11 I Tulostoma dunense I Hungary
Tulostoma dunense sp. nov.
MJ3821 I Tulostoma cf. cretaceum 4 I Hungary
FP-2023-05-11-1 I Tulostoma cf. cretaceum 3 I Kazakhstan
SNMH10 I Tulostoma cf. cretaceum 3 I Kazakhstan
FP-2023-05-11-4 I Tulostoma cf. cretaceum 3 I Kazakhstan
MJ6194 I Tulostoma cf. cretaceum 2 I Spain
MJ9304 I Tulostoma cf. cretaceum 2 I Spain
AH3955 I Tulostoma cf. cretaceum 2 I Spain
AH13672 I Tulostoma cf. cretaceum 2 I Spain
Knudsen0107 I Tulostoma cf. cretaceum 1 I Russia
NY737977 I Tulostoma cretaceum I United States I holotype
NY834492 I Tulostoma laceratum I United States
MJ7803 I Tulostoma pannonicum I Hungary
MJ7764 I Tulostoma pannonicum I Hungary I holotype
AH11603 I Tulostoma pseudopulchellum I Spain
AH11605 I Tulostoma pseudopulchellum I Spain I holotype
MJ5004 I Tulostoma sp. I Spain
MJ5996 I Tulostoma cf. kotlabae I Hungary
Finy1 I Tulostoma cf. kotlabae I Hungary
PRM 704203 I Tulostoma kotlabae I Slovakia I holotype
Bruzek140118 I Tulostoma kotlabae I Slovakia
MJ7795 I Tulostoma cf. kotlabae I Hungary
MJ4976 I Tulostoma cf. nanum I Hungary
BPI 749235 I Tulostoma lysocephalum I United States I holotype
AH15040 I Tulostoma cf. caespitosum I Spain
MJ881114 I Tulostoma cf. caespitosum I Spain
SNMH9 I Tulostoma cf. caespitosum I Slovakia
MJ3813 I Tulostoma sp. I Hungary
LISU-MGA-8 I Tulostoma lusitanicum I Portugal
Lahti201210 I Tulostoma lloydii I Italy
Kabat970428 I Tulostoma fulvellum I Slovakia
MJ8907 I Tulostoma grandisporum I Hungary I holotype
Finy10 I Tulostoma grandisporum I Hungary
FP-2017-12-09 I Tulostoma shaihuludii I Hungary
FP-2020-12-01 I Tulostoma shaihuludii I Hungary
BP112643 I Tulostoma shaihuludii I Hungary I holotype
FP-2020-12-27 I Tulostoma shaihuludii I Hungary
Tulostoma shaihuludii sp. nov.
MJ7762 I Tulostoma sp. I Hungary
MJ8701 I Tulostoma cf. fimbriatum I Spain
GB, MJ 7472 I Tulostoma fimbriatum var. punctatum I Slovakia
BPI 729033 I Tulostoma fimbriatum var. punctatum I United States I lectotype
MJ8862 I Tulostoma cyclophorum I Hungary
Cooke 2715 (NY) I Tulostoma obesum I United States I isotype
MJ8695 I Tulostoma obesum I Spain
CCB142 I Tulostoma verrucosum I United States
EL260_06 I Tulostoma squamosum I France
MJ4945 I Tulostoma subsquamosum I Hungary
HUP SH-33b I Tulostoma ahmadii I Pakistan I holotype
MLHC24 (CORD) I Tulostoma domingueziae I Argentina I holotype
BPI 704578 I Tulostoma rufum I United States I lectotype
MJ6965 I Tulostoma calcareum I Sweden I holotype
Finy4O I Tulostoma calcareum I Hungary
MJ090418 I Tulostoma melanocyclum I Hungary
Finy8 I Tulostoma fimbriatum I Hungary
GB, Manson991010 I Tulostoma fimbriatum I Sweden I epitype
JLH MyCoPortal 6604754 I Tulostoma berkeleyi I United States
MJ7761 I Tulostoma winterhoffii I Hungary
GB, MJ 8773 I Tulostoma calongei I Spain I holotype
Bruzek131207 I Tulostoma cf. pygmaeum I Slovakia
FP-2021-02-19 I Tulostoma hungaricum I Hungary
BP112641 I Tulostoma hungaricum I Hungary I holotype
FP-2019-11-23 I Tulostoma hungaricum I Hungary
FP-2022-01-02-2 I Tulostoma hungaricum I Hungary
Tulostoma hungaricum sp. nov.
AH15132 I Tulostoma submembranaceum I Mexico I holotype
FP-2021-01-24b I Tulostoma saccharioliens I Hungary
BP112642 I Tulostoma saccharioliens I Hungary I holotype
FP-2021-02-18 I Tulostoma saccharioliens I Hungary
FP-2019-12-06 I Tulostoma saccharioliens I Hungary
Tulostoma saccharioliens sp. nov.
MJ9296 I Tulostoma cf. submembranaceum I Spain
B2092 I Tulostoma albicans I Australia
MJ3844 I Tulostoma simulans I Hungary
Finy9 I Tulostoma brumale I Hungary
BPI 729284 I Tulostoma excentricum I United States I holotype
Cope NY I Tulostoma albicans I United States
Herb.Bresadola (S) I Tulostoma beccarianum I Italy I holotype
Finy2 I Tulostoma beccarianum I Hungary
AH 11698 I Tulostoma sp. I Spain
Sivertsen930717, TRH9565 I Tulostoma eckbladii I Norway I holotype
MJ8706 I Tulostoma giovanellae I Spain
MJ5015 I Tulostoma sp. I Spain
MJ7692 I Tulostoma niveum I Sweden
BPI 802484 I Tulostoma xerophilum I United States I holotype
MJ7833 I Tulostoma pulchellum I Hungary
Fritz I Tulostoma striatum I Mongolia
0.02

Figure 2. Maximum Likelihood (RAxML) tree of concatenated nrDNA ITS, nrDNA LSU and *tef1-a* sequences of representative species of the genus *Tulostoma* and the four newly-introduced species in the present study. Sequences obtained in this study are shown in bold blue. After the voucher number, the species and the country of origin are shown. Then, the type specimens are indicated. ML bootstrap support values (≥ 70) are shown before slashes and Bayesian posterior probabilities (≥ 0.90) are shown after slashes. Highlighted sections indicate affiliations to the four novel *Tulostoma* species: *T. dunense*, *T. hungaricum*, *T. sacchariolens* and *T. shaihuludii*. The illustrations exhibit basidiomata and basidiospore characteristics of the novel species. *Tulostoma pulchellum* (MJ7833) and *T. striatum* (Fritz) served as multiple outgroups. The scale bar indicates expected changes per site per branch.



Figure 3. *Tulostoma dunense*: **a, c–f** FP-2020-10-18 (BP112640, holotype), Németkér **b** FP-2021-01-02, Tázlár **a–c** basidiocarps **d, e** basidiospores **f** capillitium with basidiospores. Scale bars: 1 µm (**d**); 10 µm (**e**); 10 µm (**f**). Photos: **a, b, e, f** P. Finy **c** L. Albert **d** K. Bóka.

urban habitats, was considered a result of the maturation process. Later (Hollós 1913) corrected his earlier concept and concluded that some of the synonyms he had listed under *T. volvulatum* in his publication (Hollós 1904), in fact belonged to a complex of several species. He added illustrations of their different types of capillitia (Hollós 1913: tables 3 and 4) and concluded that his concept of *T. volvulatum* from the sand dunes was a synonym of *T. kansense*, a smooth-spored species with brownish capillitium described from North America. The capillitium in the samples from urban habitats clearly showed the undulating inner walls of the capillitium typical of *T. giovanellae* (Hollós 1913: table 3, fig. 6), although Hollós did not identify them under this name. Some 50 years later, Nagy and Babos (1969) recorded *T. giovanellae* growing on a pavement at the base of a house wall in Budapest. It matched partly the material cited by Hollós (1904) as *T. volvulatum*, but was decidedly different from the species growing in the sand dunes. Babos (1999) later came to the same conclusion. Altés et al. (1999) studied the holotype material of *T. volvulatum* and concluded that it was a synonym of *T. giovanellae* characterised by ornamented spores. They also studied the holotype material of *T. obesum* with which they identified European collections with completely smooth spores from steppe habitats and the name *T. volvulatum* was rejected. Rimóczi et al. (2011) accordingly identified the Hungarian species of the sand dunes as *T. obesum*. Molecular data (Jeppson et al. 2017) later showed that the Hungarian "*T. obesum*" was not identical with the American holotype of *T. obesum*, but was closely related to another American species described as *T. cretaceum*. It was recovered as *T.* aff. *cretaceum* by Jeppson et al. (2017). The *T.* aff. *cretaceum* from Hungary belongs to a complex

of cryptic species with a strong geographical isolation. The type of *T. cretaceum* was studied and successfully sequenced by Gube (2009), but the ITS and LSU sequences have remained unpublished. We have kindly received these sequences from Matthias Gube allowing us to include them in the phylogenetic analyses. The phylogenetic analyses showed that the type of *T. cretaceum* formed a distinct lineage within this complex (Fig. 1), proving that this North American species is different from the European and Asian lookalikes. Therefore, the Hungarian collections are proposed here as a novel species, *T. dunense*, which is closely related to samples of *T.* aff. *cretaceum* collected in Hungary, Kazakhstan and in the Russian Federation as well as in Spain (Fig. 1). The main features to distinguish *T. dunense* from the other species in the complex are mainly phylogenetic- and geographical-based data. *Tulostoma dunense* has been a protected species under Hungarian law since 2005, but to date, it has erroneously been treated under various misinterpreted and dubious names, i.e. *T. volvulatum*, *T. obesum* and *T.* aff. *cretaceum*. The ITS region of *T. dunense* differs from its closest clade represented by a single sequence (*T.* cf. *cretaceum* MJ3821, see Fig. 2) by at least 13 substitution and indel positions, which is a similarity of 98%. This sequence might represent a different species, but further collections need to be studied to clarify its taxonomic status. In contrast, low intraspecific genetic variation was detected in *T. dunense* (0–4 substitution and indel positions). The ITS and LSU sequences of an old collection identified by Long (www.mycoportal.org) under the name *Schizostoma laceratum* (NY834492) collected in 1941 in New Mexico, were provided for us by Matthias Gube. Our phylogenetic analyses indicate that this specimen belongs to the *T. cretaceum* complex as a distinct lineage. On the other hand, the nomenclature of the genus *Schizostoma*, as well as the species *S. laceratum* (Fries 1829; Léveillé 1846), seems to be problematic and needs further clarification.

Specimens examined. HUNGARY, Bács-Kiskun, Ágasegyháza, in open sand, 18 Feb 2021, P. Finy, FP-2021-02-18 (ELTE); Bócsa, in open sand, 7 Dec 2019, P. Finy, FP-2019-12-07 (ELTE); Fülöpháza, 11 Apr 2006, T. Knutsson, T. Gunnarsson, J. Jeppson, M. Jeppson, MJ7759 (GB), Ibidem, in open sand, 5 Jun 2016, P. Finy, FP-2016-06-05 (ELTE); Izsák (Soltszentimre), in open sand, 21 Nov 2019, A. Nagy, B. Dima, DB-2021-11-21-2 (ELTE); Kéleshalom, in open sand, 6 Dec 2015, P. Finy, FP-2015-12-06 (ELTE); Ibidem, in open sand, 2 Jan 2022, P. Finy, I. Ölvedi, FP-2022-01-02-1 (ELTE); Tázlár, in open sand, 11 Dec 2016, P. Finy, FP-2016-12-11 (ELTE); Ibidem, in open sand, 2 Jan 2021, P. Finy, FP-2021-01-02 (ELTE). Pest, Örkény, former military training field, sand steppe vegetation, in open sand, 5 Nov 2001, J. Jeppson, M. Jeppson, MJ6103 (GB), Ibidem, in open sand, 6 Dec 2020, P. Finy, L. Albert, FP-2020-12-06 (ELTE).

***Tulostoma hungaricum* Finy, Jeppson, L. Albert, Ölvedi & Dima, sp. nov.**

MycoBank No: MB 849932

Fig. 4

Holotype. HUNGARY, Bács-Kiskun, Bócsa, open sandy grassland, on sandy sites with scattered vegetation, near *Juniperus communis* shrubs 24 Jan 2021, P. Finy, FP-2021-01-24a (BP112641, isotype GB). GenBank: ITS OR722630, LSU OR722653.

Etymology. With reference to Hungary where it was discovered.

Description. Spore-sac subglobose, 3–6 mm. Exoperidium hyphal, heavily encrusting sand grains. Endoperidium white, pitted from adhering sand grains. Mouth small, fibrillose- fimbriate with a small and inconspicuous mouth. Socket inconspicuous. Stem slender, 9–15 × 1–1.5 mm, whitish, not bulbous. Gleba ochraceous brown. Capillitium elastic, 2–6 µm in diameter with walls 0.5–2 µm in diameter and moderate branching. Septa in general not widened. Basidiospores subglobose, 4.9–5.7 × 4.5–5.1 µm (av. 5.2 × 4.8 µm), varied in size, with fine, but visible ornamentation. SEM-photos show low verrucae coalescing in short lines.

Habitat and distribution. *Tulostoma hungaricum* occurs in the calcareous, sandy steppe areas, in dry and exposed habitats on bare sand. It has, to date, been found on the sheltered and sun-exposed, extremely warm sandy spots on the south-facing sides of *Juniperus communis*. So far, it has only been found in few localities of the Kiskunság National Park, Central Hungary.

Notes. *Tulostoma hungaricum* is the smallest *Tulostoma* species in Europe. It sometimes shares its habitat with *T. pannonicum*, another species forming small basidiomata. The latter is, however, easily distinguished on its ochraceous stem, membranous exoperidium and smaller spores. *Tulostoma hungaricum* is an isolated species belonging to the well-supported Clade 7 according to Jeppson et al. (2017), together with *T. submembranaceum* from Mexico, *T.* cf. *submembranaceum* from Spain and the below-described new species *T. sacchariolens*. In the ITS region, *T. hungaricum* differs from its closest species (*T. submembranaceum*, see Fig. 1) by almost 90 substitution and indel positions, which is a similarity of 87%. Low intraspecific genetic variability was observed in *T. hungaricum* by a difference of 0–3 substitution and indel positions.



Figure 4. *Tulostoma hungaricum*: **a, c–g** FP-2021-01-24a (BP112641, holotype), Bócsa **b** FP-2019-11-23, Orgovány **a–c** basidiocarps **d, e** basidiospores **f, g** capillitium with basidiospores. Scale bars: 1 µm (**d**); 10 µm (**e**); 20 µm (**f, g**). Photos: **a, b, e–g** P. Finy **c** L. Albert **d** K. Bóka.

Specimens examined. HUNGARY, Bács-Kiskun, Kéleshalom, open sandy grassland, near *Juniperus communis*, 2 Jan 2022, P. Finy, I. Ölvedi, L. Albert, FP-2022-01-02-2 (ELTE); Orgovány, open sandy grassland, near *Juniperus communis*, 23 Nov 2019, P. Finy, I. Ölvedi, L. Albert, FP-2019-11-23 (ELTE); Ibidem, open sandy grassland, near *Juniperus communis*, 19 Feb 2021, P. Finy, I. Ölvedi, L. Albert, FP-2021-02-19 (ELTE).

Morphologically examined specimens. HUNGARY, Bács-Kiskun, Bócsa, open sandy grassland, near *Juniperus communis*, 3 Dec 2022, P. Finy, I. Ölvedi, L. Albert, FP-2022-12-03 (herb. Finy); Fülöpháza, open sandy grassland, near *Juniperus communis*, 14 Jan 2023, P. Finy, I. Ölvedi, L. Albert, FP-2023-01-14 (herb. Finy); Pest, Tatárszentgyörgy, open sandy grassland, near *Juniperus communis*, 17 Dec 2022, I. Ölvedi, OP-2022-12-17 (herb. Ölvedi).

Tulostoma sacchariolens **Finy, Jeppson, L. Albert, Ölvedi & Dima, sp. nov.**

MycoBank No: MB 849933

Fig. 5

Holotype. HUNGARY, Bács-Kiskun, Bócsa, open disturbed sandy grassland, in a sand pit, on bare ground, 24 Jan 2021, I. Ölvedi, P. Finy, L. Albert, OP20210124 (BP112642, isotype GB). GenBank: ITS OR722632, LSU OR722654, tef1 OR707020.

Etymology. The epithet refers to its unique sweetish floral smell reminiscent of that of, for example, *Hebeloma sacchariolens*.

Description. Spore-sac subglobose, often flattened to depressed or hemispherical, 5–9 mm. Exoperidium hyphal, heavily encrusting sand, more persistent at the base of the spore-sac. Endoperidium white or dirty white, pitted from detached sand grains. Mouth delicately fimbriate. Socket conspicuous, forming a thickening on the upper part of the stem. Stem 25–50 × 1.5–2.5 mm, whitish, ornamented with orange to reddish fibrils, with age remarkably blackening, thickening towards the base, bulbous. The mature basidiomata have a pronounced sweet floral smell, reminiscent of *Hebeloma sacchariolens* Quél. or *Freesia* flowers. Gleba ferruginous brown. Capillitium 2.5–7 µm in diameter with walls 0.8–2.2 µm in diameter, lumen in general scarce, mostly straight, little branching. Most septa slightly widened. Spores subglobose, 4.6–5.3 × 4.1–5 µm (av. 4.6–4.9 µm), with coarse elongated ornamentation. SEM-photos show developed crests arranged in lines.

Habitat and distribution. Recorded in calcareous, sandy steppe areas, mostly in sunny open habitats with sparse vegetation, often in trampled or otherwise disturbed places. It is currently known only from a few localities in the sandy areas of the Danube–Tisza interfluves in Central Hungary.

Notes. With its fragrant smell and blackening stem, *Tulostoma sacchariolens* has a unique combination of characters within the genus, easily separating it from any known *Tulostoma* species. *Tulostoma sacchariolens* belongs to Clade 7 according to Jeppson et al. (2017) together with *T.* cf. *submembranaceum* (MJ9296, see Fig. 2) from Spain, *T. submembranaceum* from Mexico and the above-described *T. hungaricum*. It differs from its sister species (*T.* cf. *submembranaceum*) in the ITS region by more than 20 substitution and indel positions, which is a similarity of 96%. The intraspecific genetic variability in the ITS region amongst three sequences of *T. sacchariolens* was zero (Fig. 1), while the ITS sequence of FP-2019-12-06 had six polymorphic sites.



Figure 5. *Tulostoma sacchariolens*: **a, e** FP-2021-02-18, Orgovány **b** FP-2019-12-06, Bócsa **c, d, f** OP-2021-01-24 (BP112642, holotype), Bócsa **a–c** basidiocarps **d, e** basidiospores **f** capillitium with basidiospores. Scale bars: 1 µm (**d**); 10 µm (**e**); 20 µm (**f**). Photos: **a, b, e, f** P. Finy **c** L. Albert **d** K. Bóka.

Specimens examined. HUNGARY, Bács-Kiskun: Bócsa, open sandy grassland, 6 Dec 2019, P. Finy, FP-2019-12-06 (ELTE); Ibidem, open sandy grassland, 24 Jan 2021, P. Finy, I. Ölvedi, L. Albert, FP-2021-01-24b (ELTE); Orgovány, open sandy grassland, 18 Feb 2021, P. Finy, I. Ölvedi FP-2021-02-18 (ELTE).

Morphologically examined specimens. HUNGARY, Bács-Kiskun: Bócsa, open sandy grassland, 3 Dec 2022, P. Finy, FP-2022-12-03 (herb. Finy); Fülöpháza, open sandy grassland, 14 Jan 2023, P. Finy, FP-2023-01-14 (herb. Finy); Orgovány, open sandy grassland, 4 Dec 2021, P. Finy, I. Ölvedi, L. Albert, FP-2021-12-04 (herb. Finy); Pest, Örkény, open sandy grassland, 12 Jan 2022, I. Ölvedi, OP-2022-01-12 (herb. Ölvedi); Ibidem, open sandy grassland, 10 Dec 2022, P. Finy, I. Ölvedi, L. Albert, FP-2022-12-10 (herb. Finy).

Tulostoma shaihuludii Finy, Jeppson, L. Albert, Ölvedi, D.G. Knapp & Dima, sp. nov.

MycoBank No: MB 849934
Fig. 6

Holotype. HUNGARY, Bács-Kiskun, Tázlár, open sandy grassland, 11 Dec 2016, P. Finy, FP-2016-12-11 (BP112643, isotype GB). GenBank: ITS OR722637, LSU OR722657, tef1 OR707019.

Etymology. The epithet refers to its being reminiscent of the sandworm Shai-Hulud of the fictional planet Arrakis from the science fiction novel series Dune by Frank Herbert.

Description. Spore-sac subglobose, often flattened to depressed, 7–18 mm, relatively small compared to the size of the stem. Exoperidium hyphal, encrusting sand at the base of the spore-sac. Endoperidium white or greyish-white, pitted from detached sand grains. Mouth fimbriate, somewhat prominent. Socket developed, slightly separated from the stem. The spore-sac rarely detaches from the stem. Stem 30–70 × 3–6 mm, yellowish-brown to orange brown or reddish-brown, with age darkening, longitudinally furrowed, scaly, often curved, at the base slightly bulbous, with a conspicuous, but fragile pseudorhiza. Gleba ferruginous-cinnamon-brown. Capillitium 3.5–7 µm in diameter with walls 0.3–3.2 µm in diameter, mostly straight, sparsely branched, inner wall often undulating. Septa not or slightly widened. Abundant, thin-walled, septate capillitium hyphae present amongst normal capillitium threads. Basidiospores globose, sometimes flattened, 4.1–5.2 × 3.5–4.7 µm (av. 4.1 × 4.6 µm), finely asperulate, ornamentation not always visible under LM. SEM photos show fine warts arranged in lines forming a dense network.

Habitat and distribution. Occurs in dry and loose calcareous, open sandy habitats of the *Festucetum vaginatae* natural grasslands. It mainly grows solitary, deeply rooted in the sand in spots with bare sand. It is currently known only from the sandy areas of Central Hungary.

Notes. *Tulostoma shaihuludii* is similar in stature to *T. fimbriatum* and *T. winterhoffii*, but can be easily distinguished by its habitat (open sand) and its microcharacters, particularly the spore wall ornamentation. It belongs to Clade 2 according to Jeppson et al. (2017) and it forms a sister clade of *Tulostoma* cf. *fimbriatum*

Figure 6. *Tulostoma shaihuludii*: **a** FP-2020-12-01-3, Fülöpháza **b, d–g** FP-2016-12-11 (BP112643, holotype), Tázlár **c** AL-2021-01-24, Bócsa **h** FP-2017-12-09, Orgovány **a–c** basidiocarps **d, e** basidiospores **f, g** capillitium with basidiospores **h** thin-walled, septate capillitium hyphae. Scale bars: 1 µm (**d**); 10 µm (**e**); 20 µm (**f–h**). Photos: **a, b, e–h** P. Finy **c** L. Albert **d** K. Bóka.

(MJ8701 as "*T.* sp2" in Jeppson et al. (2017)) from which it differs in the ITS region by 45 substitution and indel positions, which is a similarity of 93%. The intraspecific genetic variability of *T. shaihuludii* is low (0–3 substitution and indels positions).

Specimens examined. Hungary, Bács-Kiskun, Fülöpháza, Fülöpházi homokbuckák, sand steppe vegetation, 11 Apr 2006, J. Jeppson, M. Jeppson, MJ7762 (GB); Ibidem, open sandy grassland, 1 Dec 2020, P. Finy, I. Ölvedi, FP-2020-12-01-3 (ELTE); Orgovány, open sandy grassland, 9 Dec 2017, P. Finy, FP-2017-12-09 (ELTE); Pirtó, open sandy grassland, 27 Dec 2020, P. Finy, L. Albert, FP-2020-12-27 (ELTE).

Morphologically examined specimens. Hungary, Bács-Kiskun, Bócsa, open sandy grassland, 7 Dec 2019, P. Finy, FP-2019-12-07 (herb. Finy); Ibidem, open sandy grassland, 24 Jan 2021, P. Finy, L. Albert, I. Ölvedi, FP-2021-01-24 (herb. Finy), AL-2021-01-24 (herb. Albert); Ibidem, open sandy grassland, 4 Dec 2021, P. Finy, I. Ölvedi, FP-2021-12-04 (herb. Finy); Ibidem, open sandy grassland, 3 Dec 2022, P. Finy, FP-2022-12-03 (herb. Finy); Fülöpháza, open sandy grassland, 2 Dec 2018, P. Finy, FP-2018-12-02 (herb. Finy); Ibidem, open sandy grassland, 16 Jan 2022, P. Finy, I. Ölvedi, FP-2022-01-16 (herb. Finy); Izsák (Soltszentimre), open sandy grassland, 4 Feb 2016, P. Finy, FP-2016-02-04 (herb. Finy); Ibidem, open sandy grassland, 14 Dec 2016, P. Finy, FP-2016-12-14 (herb. Finy); Ibidem, open sandy grassland, 17 Jan 2019, P. Finy, FP-2019-01-17 (herb. Finy); Ibidem, open sandy grassland, 16 Dec 2020, P. Finy, FP-2020-12-16-1 (herb. Finy); Kéleshalom, open sandy grassland, 6 Dec 2015, P. Finy, FP20151206 (herb. Finy); Ibidem, open sandy grassland, 2 Jan 2022, P. Finy, I. Ölvedi, FP-2022-01-02-3 (herb. Finy); Kiskunhalas, open sandy grassland, 22 Dec 2019, P. Finy, FP-2019-12-22 (herb. Finy); Ibidem, open sandy grassland, 5 Jan 2023, P. Finy, I. Ölvedi, FP-2023-01-05 (herb. Finy); Kunbaracs, open sandy grassland, 5 Feb 2022, P. Finy, I. Ölvedi, FP-2022-02-05 (herb. Finy); Orgovány, open sandy grassland, 13 Aug 2017, P. Finy, FP-2017-08-13 (herb. Finy); Ibidem, open sandy grassland, 18 Feb 2021, P. Finy, I. Ölvedi, FP-2021-02-18 (herb. Finy); Pirtó, open sandy grassland, 16 Jan 2016, P. Finy, FP-2016-01-16 (herb. Finy); Tázlár, open sandy grassland, 11 Dec 2016, P. Finy, FP-2016-12-11 (herb. Finy); Pest, Örkény, open sandy grassland, 12 Jan 2022, I. Ölvedi, OP-2022-01-12 (herb. Ölvedi); Tatárszentgyörgy, open sandy grassland, 1 Jan 2022, I. Ölvedi, OP-2022-01-01 (herb. Ölvedi); Ibidem, open sandy grassland, 10 Dec 2022, P. Finy, FP-2022-12-10 (herb. Finy); Ibidem, open sandy grassland, 17 Dec 2022, I. Ölvedi, OP-2022-12-17 (herb. Ölvedi); Tolna, Paks, open sandy grassland, 4 Feb 2018, P. Finy, FP-2018-02-04 (herb. Finy); Ibidem, open sandy grassland, 22 Jan 2021, P. Finy, FP-2021-01-22 (herb. Finy); Ibidem, open sandy grassland, 9 Jan 2022, I. Ölvedi, P. Finy, OP-2022-01-09 (herb. Ölvedi); Ibidem, open sandy grassland, 27 Feb 2022, P. Finy, FP-2022-02-27 (herb. Finy).

Discussion

The results of our study further emphasise the high species diversity amongst the stalked puffballs (*Tulostoma*) in East Central Europe, as previously indicated by Jeppson et al. (2017). In Hungary, so far 19 species have been recorded, including the four new species proposed in this study. The Pannonian, dry and sandy grasslands between the rivers Danube and Tisza, as well as adjacent areas in Central Hungary, harbour to date 66% of all described species of *Tulostoma* known to occur in Europe (29 spp.). The dry, sandy grasslands in Central Hungary have a long continuity as natural grasslands or as sheep pastures and

are characterised by steppe flora and fauna. Both natural and grazed habitats are rich in gasteroid fungi, but usually their species composition is different. The summer and autumn temperatures in the sand are extremely high and the yearly precipitation is low. The dry and drought-resisting basidiomata of *Tulostoma* species could be considered as adaptations to xeric conditions. The development of the basidiomata occurs mainly in late autumn and early winter.

Tulostoma species are generally rare (although locally abundant) and the current knowledge of their population structures in Europe is limited. However, their occurrences are highly dependent on the habitat status where they grow and changes in land management are likely to be detrimental to their populations. A vast majority of the European *Tulostoma* species are Red-Listed in the countries where they occur (http://www.eccf.eu/redlists-en.ehtml).

In addition to the four novel species proposed herein, the results from previous works (e.g. Jeppson et al. (2017)) and our ongoing studies indicate the presence of many more undescribed species of *Tulostoma* in Central Europe.

Acknowledgements

The work was supported by the ÚNKP-22-5 New National Excellence Program of the Ministry for Culture and Innovation from the source of the National Research, Development and Innovation Fund, the National Research, Development and Innovation Office of Hungary (FK-143061, the ELTE Institutional Excellence Program 2020 (TKP2020-IKA-05), Diagnostics and Therapy 2). Bálint Dima is grateful to the János Bolyai Research Scholarship of the Hungarian Academy of Sciences. We thank to Csilla Gergely for her assistance in laboratory work. We are grateful to Matthias Gube (Germany) for allowing us to publish the type sequences of *Tulostoma cretaceum* and the sequences of *Schizostoma laceratum*.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

New National Excellence Program of the Ministry for Culture and Innovation from the source of the National Research, Development and Innovation Fund (Hungary); National Research, Development and Innovation Office (Hungary); János Bolyai Research Scholarship of the Hungarian Academy of Sciences.

Author contributions

Conceptualisation: PF, BD. Methodology: PF, MJ, DGK, VP, KB, DV, BD. Validation: PF, MJ, LA, IÖ. Formal analysis: PF, MJ, DGK, DV, BD. Investigation: PF, MJ, LA, IÖ, KB, BD. Resources: PF, MJ, LA, IÖ, BD. Data Curation: PF, MJ, DGK, LA, IÖ, DV, BD. Writing - Original draft: PF, MJ, VP, BD. Writing - Review and Editing: PF, MJ, DGK, VP, LA, IÖ, KB, DV, GMK, BD. Visualisation: PF, DGK, VP, LA, IÖ, KB. Supervision: MJ, GMK, BD. Funding Acquisition: GMK, BD.

Author ORCIDs

Dániel G. Knapp https://orcid.org/0000-0002-7568-238X
Viktor Papp https://orcid.org/0000-0001-6994-8156
Károly Bóka https://orcid.org/0000-0002-1324-3592
Gábor M. Kovács https://orcid.org/0000-0001-9509-4270
Bálint Dima https://orcid.org/0000-0003-2099-3903

Data availability

All the data that support the findings of this study are available in the main text or in publicly accessible data repositories, as indicated in the text.

References

Altés A, Moreno G, Wright JE (1999) Notes on *Tulostoma volvulatum* and *T. giovanellae*. Mycological Research 103(1): 91–98. https://doi.org/10.1017/S0953756298006704

Babos M (1999) Higher fungi (Basidiomycotina) of the Kiskunság National Park and its environs. In: Lőkös L, Rajczy M (Eds) The flora of the Kiskunság National Park 2. Cryptogams, Hungarian Natural History Museum, Budapest, 199–298.

Bölöni J, Molnár Zs, Kun A [Eds] (2011) Magyarország élőhelyei. A hazai vegetációtípusok leírása és határozója. ÁNÉR 2011. MTA ÖBKI, Vácrátót.

Caffot MLH, Domínguez LS, Hosaka K, Crespo EM (2011) *Tulostoma domingueziae* sp. nov. from *Polylepis australis* woodlands in Córdoba Mountains, central Argentina. Mycologia 103(5): 1047–1054. https://doi.org/10.3852/10-266

Fries EM (1829) Systema mycologicum. Vol. 3. Gryphiswaldiae, 524 pp. + 202 pp.

Gardes M, Bruns TD (1993) ITS primers with enhanced specifity for basidiomycetes – application to the identification of mycorrhizae and rusts. Molecular Ecology 2(2): 113–118. https://doi.org/10.1111/j.1365-294X.1993.tb00005.x

Gouy M, Tannier E, Comte N, Parsons DP (2021) Seaview version 5: A multiplatform sofware for multiple sequence alignment, molecular phylogenetic analyses, and tree reconciliation. In: Katoh K. (Ed.) Multiple Sequence Alignment. Methods in Molecular Biology, vol 2231. Humana, New York, NY, 241–260. https://doi.org/10.1007/978-1-0716-1036-7_15

Gube M (2009) Ontogeny and phylogeny of gasteroid members of Agaricaceae (Basidiomycetes). PhD Dissertation, University of Jena, Jena, Germany, 145 pp. https://d-nb.info/999990691/34

Hussain S, Yousaf N, Afshan NS, Niazi AR, Ahmad H, Khalid AN (2016) *Tulostoma ahmadii* sp. nov. and *T. squamosum* from Pakistan. Turkish Journal of Botany 40: 218–225. https://doi.org/10.3906/bot-1501-9

Hollós L (1904) Die Gasteromyceten Ungarns. Oswald Weigel, Leipzig.

Hollós L (1913) Zu den Gasteromyceten Ungarns – Magyarország Gasteromycetái-hoz. Magyar Botanikai Lapok 12(6–7): 194–200.

Jeppson M, Altes A, Moreno G, Nilsson RH, Loarce Y, de Bustos A, Larsson E (2017) Unexpected high species diversity among European stalked puffballs – a contribution to the phylogeny and taxonomy of the genus *Tulostoma* (Agaricales). MycoKeys 21: 33–88. https://doi.org/10.3897/mycokeys.21.12176

Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. Molecular Biology and Evolution 30(4): 772–780. https://doi.org/10.1093/molbev/mst010

Kumar S, Stecher G, Tamura K (2016) MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets. Molecular Biology and Evolution 33(7): 1870–1874. https://doi.org/10.1093/molbev/msw054

Léveillé J-H (1846) Description des champignons de l'herbier du muséum de Paris. Annales des Sciences Naturelles sér. 3. Botanique 5: 111–166.

Moreno G, Altés A, Wright JE (1992) *Tulostoma squamosum*, *T. verrucosum* and *T. mussooriense* are the same species. Mycotaxon 43: 61–68.

Moreno G, Altés A, Ochoa C, Wright JE (1997) Notes on type materials of *Tulostoma*. Some species with mixed holotypes. Mycological Research 101(8): 957–965. https://doi.org/10.1017/S0953756297003572

Nagy L, Babos M (1969) Vorkommen einer seltenen Stielbovist-Art (*Tulostoma giovanellae* Bres. in Ungarn). Mikológiai Közlemények 1969(2): 115–121.

Papp V, Dima B (2018) New systematic position of *Aurantiporus alborubescens* (Meruliaceae, Basidiomycota), a threatened old-growth forest polypore. Mycological Progress 17(3): 319–332. https://doi.org/10.1007/s11557-017-1356-3

Persoon CH (1794) Neuer Versuch einer systematischen Eintheilung der Schwämme. Neues Magazin für die Botanik 1: 63–128.

Persoon CH (1801) Synopsis Methodica Fungorum. Gottingae.

Rehner SA, Buckley E (2005) A *Beauveria* phylogeny inferred from nuclear ITS and EF1-a sequences: Evidence for cryptic diversification and links to *Cordyceps* teleomorphs. Mycologia 97(1): 84–98. https://doi.org/10.3852/mycologia.97.1.84

Rehner SA, Samuels GJ (1994) Taxonomy and phylogeny of *Gliocladium* analysed from nuclear large subunit ribosomal DNA sequences. Mycological Research 98(6): 625–634. https://doi.org/10.1016/S0953-7562(09)80409-7

Rimóczi I, Jeppson M, Benedek L (2011) Characteristic and rare species of Gasteromycetes in Eupannonicum. Fungi Non Delineati : Raro Vel Haud Perspecte et Explorate Descripti aut Definite Picti: 56–57. [Edizioni Candusso, Alassio]

Ronquist F, Huelsenbeck JP (2003) MRBAYES 3: Bayesian phylogenetic inference under mixed models. Bioinformatics (Oxford, England) 19(12): 1572–1574. https://doi.org/10.1093/bioinformatics/btg180

Rusevska K, Calonge FD, Karadelev M, Martín MP (2019) Fungal DNA barcode (ITS nrDNA) reveals more diversity than expected in *Tulostoma* from Macedonia. Turkish Journal of Botany 43(1): 102–115. https://doi.org/10.3906/bot-1804-38

Schoch CL, Seifert KA, Huhndorf S, Robert V, Spouge JL, Levesque CA, Chen W, Bolchacova E, Voigt K, Crous PW, Miller AN, Wingfield MJ, Aime MC, An K-D, Bai F-Y, Barreto RW, Begerow D, Bergeron M-J, Blackwell M, Boekhout T, Bogale M, Boonyuen N, Burgaz AR, Buyck B, Cai L, Cai Q, Cardinali G, Chaverri P, Coppins BJ, Crespo A, Cubas P, Cummings C, Damm U, de Beer ZW, de Hoog GS, Del-Prado R, Dentinger B, Diéguez-Uribeondo J, Divakar PK, Douglas B, Dueñas M, Duong TA, Eberhardt U, Edwards JE, Elshahed MS, Fliegerova K, Furtado M, García MA, Ge Z-W, Griffith GW, Griffiths K, Groenewald JZ, Groenewald M, Grube M, Gryzenhout M, Guo L-D, Hagen F, Hambleton S, Hamelin RC, Hansen K, Harrold P, Heller G, Herrera C, Hirayama K, Hirooka Y, Ho H-M, Hoffmann K, Hofstetter V, Högnabba F, Hollingsworth PM, Hong S-B, Hosaka K, Houbraken J, Hughes K, Huhtinen S, Hyde KD, James T, Johnson EM, Johnson JE, Johnston PR, Jones EBG, Kelly LJ, Kirk PM, Knapp DG, Kõljalg U, Kovács GM, Kurtzman CP, Landvik S, Leavitt SD, Liggenstoffer AS, Liimatainen K, Lombard L, Luangsa-ard JJ, Lumbsch HT, Maganti H, Maharachchikumbura SSN, Martin MP, May TW, McTaggart AR, Methven AS, Meyer W, Moncalvo J-M, Mongkolsamrit S, Nagy LG, Nilsson RH, Niskanen T, Nyilasi I, Okada G, Okane I, Olariaga I, Otte J, Papp T, Park

D, Petkovits T, Pino-Bodas R, Quaedvlieg W, Raja HA, Redecker D, Rintoul TL, Ruibal C, Sarmiento-Ramírez JM, Schmitt I, Schüßler A, Shearer C, Sotome K, Stefani FOP, Stenroos S, Stielow B, Stockinger H, Suetrong S, Suh S-O, Sung G-H, Suzuki M, Tanaka K, Tedersoo L, Telleria MT, Tretter E, Untereiner WA, Urbina H, Vágvölgyi C, Vialle A, Vu TD, Walther G, Wang Q-M, Wang Y, Weir BS, Weiß M, White MM, Xu J, Yahr R, Yang ZL, Yurkov A, Zamora J-C, Zhang N, Zhuang W-Y, Schindel D (2012) Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi. Proceedings of the National Academy of Sciences of the United States of America 109(16): 6241–6246. https://doi.org/10.1073/pnas.1117018109

Siller I, Vasas G (1995) Red list of macrofungi of Hungary (revised edition). Studia Botanica Hungarica 26: 7–14.

Siller I, Vasas G, Pál-Fám F, Bratek Z, Zagyva I, Fodor L (2005) Hungarian distribution of the legally protected macrofungi species. Studia Botanica Hungarica 36: 131–163.

Siller I, Dima B, Albert L, Vasas G, Fodor L, Pál-Fám F, Bratek Z, Zagyva I (2006) Protected macrofungi in Hungary. Mikol. Közlemények. Clusiana 45(1–3): 3–158.

Silvestro D, Michalak I (2012) RaxmlGUI: A graphical front-end for RAxML. Organisms, Diversity & Evolution 12(4): 335–337. https://doi.org/10.1007/s13127-011-0056-0

Staden R, Beal KF, Bonfield JK (2000) The Staden package, 1998. In: Misener S, Krawetz SA (Eds) Bioinformatics methods and protocols. Humana Press, 115–130. https://doi.org/10.1385/1-59259-192-2:115

Stamatakis A (2014) RAxML Version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. Bioinformatics 30: 1312–1313. https://doi.org/10.1093/bioinformatics/btu033

Varga T, Krizsán K, Földi Cs, Dima B, Sánchez-García M, Sánchez-Ramirez S, Szőllősi G, Szarkándi GJ, Papp V, Albert L, Angelini C, Antonín V, Bougher N, Buchanan P, Buyck B, Bense V, Catcheside P, Cooper J, Dämon W, Desjardin D, Finy P, Geml J, Hughes K, Justo AF, Karasiński D, Kautmanova I, Kerr S, Kiss B, Kocsubé S, Kotiranta H, Lechner BE, Liimatainen K, Lukács Z, Morgado L, Niskanen T, Noordeloos ME, Ortiz-Santana B, Ovrebo C, Rácz N, Savchenko A, Shiryaev A, Soop K, Spirin V, Szebenyi Cs, Tomsovsky M, Tulloss RE, Uehling J, Vágvölgyi Cs, Papp T, Martin FM, Miettinen O, Hibbett DS, Nagy LG (2019) Megaphylogeny resolves global patterns of mushroom diversification. Nature Ecology & Evolution 3(4): 668–678. https://doi.org/10.1038/s41559-019-0834-1

Vilgalys R, Hester M (1990) Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. Journal of Bacteriology 172(8): 4238–4246. https://doi.org/10.1128/jb.172.8.4238-4246.1990

White TJ, Bruns T, Lee S, Taylor JW (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ (Eds) PCR protocols: A guide to the methods and applications. Academic Press, New York, 315–322. https://doi.org/10.1016/B978-0-12-372180-8.50042-1

Wright JE (1987) The genus *Tulostoma* (Gasteromycetes) – A world monograph. Bibliotheca Mycologica 113. Berlin, Stuttgart, 338 pp.